

Original Article

**IN-VITRO ANTI-CANCER STUDY OF BIOTRANSFORMED MATERIAL (EXTRACT) OF
CALENDULA OFFICINALIS BY MTT ASSAY**

PRAKASH ITANKAR*, ASHLESH MURKUTE¹

University Department of Pharmaceutical Sciences, R.T.M. Nagpur University Nagpur, Mahatma Jyotiba Phule Shaikshanik Parisar,
Amaravati road, Nagpur 440033 Maharashtra, India.
Email: prakashitankar@hotmail.com

Received: 25 Jun 2014 Revised and Accepted: 01 Aug 2014

ABSTRACT

Objective: This work aims to determine the *in-vitro* anti-cancer activity of biotransformed material (extract) of *calendula officinalis* flowers.

Methods: Various concentrations of the biotransformed material (extract) of *calendula officinalis* were tested. The MTT assay method was used using A549 cell lines, LN Cap-FGC cell lines and MDA-MB cell lines.

Results: The result shows promising effect of biotransformed material (extract) of *calendula officinalis*. The IC₅₀ values were found to be less than 10 µg against A549 cell lines, the activities against LN Cap-FGC cell lines were found to be 10 µg and IC₅₀ values were found to be 10 µg against MDA-MB cell lines

Conclusion: From this study it was evident that the cancer cell growth was significantly inhibited by treatment with biotransformed material (extract). Hence, *in vitro* studies of *Calendula officinalis* was found to have a potential anti-cancer activity.

Keywords: *Calendula officinalis*, MTT assay, A 549 cell lines, LN Cap-FGC cell lines, MDA-MB cell lines, anti-cancer activity.

INTRODUCTION

A proper health care system can be established, supplying low cost medicine to population by using various medicinal plants. Medicinal plants are usually used for Ayurvedic, Unani and other treatments in rural areas. Recent discovery shows that these plants have fewer side effects than the Allopathic medicine. The medicinal importance of these plants is due to the presence of secondary metabolites present in them. Some of the important bioactive compounds are alkaloids, glycosides, triterpenoids, terpenoids, flavonoids, polyphenols, reducing sugars, saponins, steroids and tannins etc. Among the plants, *Calendula officinalis* has been used in ayurvedic medicine and has a wide number of traditional uses including internal use for blepharitis, eczema, gastritis, minor burns including sunburns, warts and minor injuries such as sprains and wounds. Calendula flowers have been considered beneficial in reducing inflammation and wound healing. In homeopathy, Calendula ointments are also used in the healing stages of second and third degree burns to stimulate regrowth of skin and to diminish scar formation [1].

Cancer is the abnormal growth of cells with uncontrolled division resulting in increased number of cells [2]. In spite of a number of new anti cancer drugs the prevalence of cancer has been found increased now-a-days. Cancer is considered as one of the serious disease which is characterized by abnormal or uncontrolled growth of tissues. It has become a major cause of death and there is an urgent need for its control. The most important drawback of the current cancer therapeutic practices such as chemotherapy and radiation therapy is the suppression of immune system. Since plants have been a prime source of natural products for the treatment of various diseases and they are also highly effective conventional drugs for the treatment of many forms of cancer [3-5]. *Calendula officinalis* belonging to family Asteraceae has been widely used in homeopathic medicine for the treatment of many diseases [6]. It has been reported to possess many pharmacological activities, which include antioxidant [7], anti-inflammatory [8], antibacterial [9], antifungal [10] and antiviral activity [11]. It also possess cytotoxic as well as tumor reducing potential [12]. Literature survey revealed that *Calendula officinalis*, is used in traditional medicine all over the world for its antiseptic property as well as to treat certain skin disorders.

The process in which suitable modification of substrate structure is achieved by biological enzyme catalysis is termed as biotransformation.

There are the following three ways with the help of which the biotransformation process is carried out.

1. With purified culture of microorganisms.
Ex. Bacteria or Fungi or Yeast
2. With plant cell culture or tissue culture
3. With isolated purified microbial or plant or animal cell enzyme.

In the current context, general goals of biotransformation can be listed as follows:

- Specific modification of substrate molecule via selective transformation reaction
- Partial degradation of substrate molecule into desirable metabolites by means of controlled microbial reaction or reaction pathway.
- Extension of substrate or by the use of biosynthetic reactions to form artificial products.

The first processes which were successfully accomplished by biotransformation were.

- The oxidation of alcohol to acetic acid by *Bacterium xylinum*.
- The oxidation of glucose to gluconic acid by *Acetobacter acetii*.
- Acyloin formation from benzaldehyde and acetaldehyde through yeast.

But, biotransformation gained its present status only after discovery of microbial transformations of steroids viz. Reduction of androstenedione to testosterone through yeast, 11-hydroxylation of progesterone by *Rhizopus arrhizus* (Peterson & Murray 1952).

In the aftermath, diverse transformations of substrates from many different groups and types of compounds were found, some of which were useful in chemosynthesis.

Very few biotransformation studies have been carried out regarding the chemical study of biotransformed products and their pharmacological evaluation. Biotransformation with alcohol, with water have been done however the biotransformation with cow urine has not been done yet. It was therefore envisaged to evaluate the biotransformation potential of Cow urine as an extraction medium and subsequent impact on the efficacy of its extract.

Microbial biotransformation employs microorganisms for achieving the desired conversion of various substrates. Microorganisms can catalyze the reactions which otherwise are difficult to carry out. The success of microbial biotransformation depends greatly on proper selection of microorganisms. The well known enrichment techniques are used for isolation of potent microbial agents. These often employ enrichment cultures obtained from soils, decomposing materials, sewage plants, river water or factory grounds etc. The sample is incubated in a growth medium containing substrate to be transformed as sole carbon source. Microorganisms employ both constitutive as well as inducible enzymes to degrade and synthesize a great variety of compounds. The discovery of the phenomenon of microbial biotransformation of complicated organic compounds provided impetus to employ microorganisms in the production of novel agents. Gradually, the methods of preparing new derivatives of known substrates were applied to modify organic compounds like Carbohydrates, steroids, terpenoids, flavonoids, alkaloids, antibiotics, amino acids etc. Such alteration encompasses many changes in the substrate moiety by the addition, degradation or modification. When basic structure of resultant molecule remains unaltered, the microbial biotransformation can be applied for formation of desired derivatives of precursor. Traditionally, cow urine itself has some medicinal properties and employing it in the process of extraction through biotransformation is envisaged to have positive improvement in the original potency of the herb under study.

MATERIALS AND METHODS

Plant Material

Procurement of *Calendula officinalis* flowers

Controlled dried flowers of *Calendula officinalis* were procured from the horticulture field at Nagpur.

Cow urine was procured from Govigyan Anusandhan Kendra, Dewlapur, Nagpur

Authentication

The flowers were authenticated by Dr. Dongarwar, Department of Botany, R.T.M. Nagpur University, Nagpur, with the herbarium voucher specimen number 9763.

Preparation of Extract

The collected flowers about 200 gm. were macerated in cow urine (1 Lit.) for time period of 28 days [13]. After the completion of this time period, the biotransformed material (Extract) was harvested, lyophilized, powdered (20 Gm.) which was then used for further studies.

Preparation of Aqueous Extract for comparative study

The collected flowers about 200 gm. were macerated in water (1 Lit.) for time period of 28 days [13]. After the completion of this time period, the biotransformed material (Extract) was harvested, lyophilized, powdered (20 Gm.) which was then used for further studies.

Maintenance of Cell culture

The A549 cell lines, LN Cap- FGC cell lines and MDA-MB cell lines were maintained at The Maratha Mandal Dental College, Belgaum, Karnataka. Cell lines were grown in Minimal essential medium (MEM) supplemented with 4.5 g/L glucose, 2 m ML-glutamine and 5% fetal bovine serum (FBS) (growth medium) at 37 °C in 5% CO₂ incubator.

MTT Assay (MICRO TITER ASSAY)

The *in vitro* MTT assay method was used to determine the inhibitory effects of test compounds on cell growth. The trypsinized cells from T-25 flask were seeded in each well of 96-well flat-bottomed tissue culture plate at a density of 5x10³ cells/well in growth medium and cultured at 37 °C in 5% CO₂ to adhere. After 48 hr incubation, the supernatant was discarded and the cells were pre-treated with growth medium and were subsequently mixed with different concentrations of test samples (10, 20 and 30 µg) in triplicates to achieve a final volume of 100 µl and then cultured for 48 hr. The compound was prepared as 1.0 mg/ml concentration stock solutions in PBS. Culture medium and solvent were used as control and blank respectively. Each well then received 5 µl of fresh MTT (5mg/ml in PBS) followed by incubation for 2 hr at 37 °C. The supernatant growth medium was removed from the wells and replaced with 100 µl of DMSO to solubilise the colored Formosan product. After 30 min incubation, the absorbance (OD) of the culture plate was read at a wavelength of 492 nm on an ELISA reader [14].

The percentage inhibition of cancer cell lines was calculated with the following formula

$$\% \text{ inhibition} = 100 - \frac{(At - Ab)}{(Ac - Ab)} \times 100$$

Where, At: Absorbance of test,

Ab: Absorbance of blank,

Ac: Absorbance of control.

RESULTS AND DISCUSSION

The effect of biotransformed material (extract) of *Calendula officinalis* was studied using MTT assay at a dose of 10, 20, 30 µg and Vincristine was used as a standard at a dose of 90 µg indicate the efficacy of microbial biotransformed material against cancerous cells.

Hence by

Cytotoxic properties of biotransformed material (extract) of *Calendula officinalis* against A549 cell lines

The IC₅₀ value of *Calendula officinalis* against A549 cell lines was found to be Less than 10 µg and the results are provided in Table 1.

Cytotoxic properties of biotransformed material (extract) of *Calendula officinalis* against LN Cap-FGC cell lines

The IC₅₀ value of *Calendula officinalis* against LN Cap-FGC cell lines was found to be 10 µg and the results are provided in Table 2.

Cytotoxic properties of biotransformation material (extract) of *Calendula officinalis* against MDA-MB cell lines

The IC₅₀ value of *Calendula officinalis* against MDA-MB cell lines was found to be

10 µg and the results are provided in Table 3.

Table 1: Cytotoxicity effect against A549 cell lines

S. No.	Name of drug	Concentration (µg)	O.D. at 492 nm	%of cell lysis	IC ₅₀
1.	<i>Calendula officinalis</i> Biotransformation material (extract)	10	0.719	75%	<10 µg
		20	0.782	>75%	
		30	1.226	100%	
2.	Control	-	0.240	No lysis	-
3.	Vincristine	90	1.117	90%	-

Table 2: Cytotoxicity effect against LN Cap-FGC cell lines (LN Cap-FGC - Human carcinoma Prostate)

S. No.	Name of drug	Concentration (µg)	O.D. at 492 nm	%of cell lysis	IC ₅₀
1.	<i>Calendula officinalis</i> Biotransformation material (extract)	10	0.578	50%	10 µg
		20	0.768	75%	
		30	1.084	100%	
2.	Control	-	0.321	No lysis	-
3.	Vincristine	90	1.117	90%	-

Table 3: Cytotoxicity effect against MDA-MB cell lines

S. No.	Name of drug	Concentration (µg)	O.D. at 492 nm	%of cell lysis	IC ₅₀
1.	<i>Calendula officinalis</i> Biotransformation material (extract)	10	0.622	50%	10 µg
		20	0.764	75%	
		30	0.940	100%	
2.	Control	-	0.334	No lysis	-
3.	Vincristine	90	1.117	90%	-

Comparative results for Aqueous Extracts

Table 1: Cytotoxicity effect against A549 cell lines

S. No.	Name of drug	Concentration (µg)	O.D. at 492 nm	%of cell lysis	IC ₅₀
1.	<i>Calendula officinalis</i> Aqueous extract	10	0.719	No Lysis	-----
		20	0.782	No Lysis	
		30	1.226	No Lysis	
2.	Control	-	0.240	No lysis	-
3.	Vincristine	90	1.117	90%	-

Table 2: Cytotoxicity effect against LN Cap-FGC cell lines (LN Cap-FGC - Human carcinoma Prostate)

S. No.	Name of drug	Concentration (µg)	O.D. at 492 nm	%of cell lysis	IC ₅₀
1.	<i>Calendula officinalis</i> Aqueous extract	10	0.578	No lysis	-----
		20	0.768	No lysis	
		30	1.084	No lysis	
2.	Control	-	0.321	No lysis	-
3.	Vincristine	90	1.117	90%	-

Table 3: Cytotoxicity effect against MDA-MB cell lines

S. No.	Name of drug	Concentration (µg)	O.D. at 492 nm	%of cell lysis	IC ₅₀
1.	<i>Calendula officinalis</i> Aqueous extract	10	0.622	No lysis	-----
		20	0.764	No lysis	
		30	0.940	No lysis	
2.	Control	-	0.334	No lysis	-
3.	Vincristine	90	1.117	90%	-

CONCLUSION

The anti-cancer studies of the extract of *Calendula officinalis* obtained after biotransformation using cow urine as an extraction medium on A549 cell lines, LN Cap-FGC cell lines and MDA-MB cell lines was carried out by using MTT cell growth inhibition assay. The results showed that the maximum percentage inhibition of cancer cell lines by *Calendula officinalis* biotransformed extract was found to be 50% at a dose of less than 10 µg for A549 cell lines, 50% at a dose of 10 µg for LNCap-FGC cell lines and 50% at a dose of 10 µg for MDA-MB cell lines.

Hence biotransformed material (extract) of *Calendula officinalis* obtained by using cow urine as medium for extraction can be used as a potent anti-cancer agent whereas the aqueous extract of same plant material doesn't show any activity.

CONFLICT OF INTERESTS

Declared None

ACKNOWLEDGEMENTS

We wish to thank the Govigyan Anusandhan Kendra, Dewlapur, Nagpur, for providing fresh cow urine and Maratha Mandal Dental college, Belgaum, Karnataka, who gave support to carry the *in vitro* anti-cancer studies.

REFERENCES

1. Akihisa T. Triterpene alcohols from the flowers of compositae and their anti-inflammatory effects. J Phytochemistry 1996; 43(6):1225-60.
2. Manoharan et al. In Vivo and In vitro antitumor activity of *Jasminum sambu* (linn) AIT Oleaceae flower against Dalton's ascites lymphoma induced Swiss albino mice. Int J Pharm Pharm Sci 2012;4(1):144-7.
3. Mangale SM, Chonde SG, Raut PD. Use of *Moringaoleifera* (drumstick) seed as natural absorbent and an antimicrobial agent for ground water treatment. Res J Rec Sci 2012;1(3):31-40.

4. Mondal D, Mondal T. A review on efficacy of *Azadirachta indica* A. juss based biopesticides: an Indian perspective. Res J Rec Sci 2012;1(3):94-9.
5. Jocelyn PC. The effect of glutathione on protein sulphydryl groups in Rat-liver homogenates. J Biochem 1962;85:480-5.
6. Zitterl-Eglseer K, Sosa S, Jurenitsch J, Schubert-Zsilavecz M, Della Loggia R, Tubaro A, et al. Anti-edematous activities of the main triterpendiol esters of marigold (*Calendula officinalis* L.) J Ethnopharmacol 1997;57:139-44.
7. Preethi KC, Kuttan G, Kuttan R. Antioxidant potential of *Calendula officinalis* flowers *in vitro* and *in vivo*. J Pharm Biol 2006;44(9):691-7.
8. Della Loggia R, Tubaro A, Sosa S, Becker H, Saar S, Issac O. The role of triterpenoids in the topical anti-inflammatory activity of *Calendula* flowers. J Planta Med 1994;60:516-20.
9. Dumenil G, Chemli R, Balausard G. Evaluation of antibacterial properties of *Calendula officinalis* flowers and mother homeopathic tinctures of *Calendula officinalis*. J Ann Pharma Fran 1980;38:493-9.
10. Kasiram K, Sakharkar PR, Patil AT. Antifungal activity of *Calendula officinalis*. Indian J Pharm Sci 2000;6:464-6.
11. Barbour EK, Sagherian V, Talhouk S, Talhouk R, Farran MT, Sleiman FT, Harakeh S. Evaluation of homeopathy in broiler chickens exposed to live viral vaccines and administered *Calendula officinalis* extract. J Med Sci Monit 2004;10(8):281-5.
12. Boucaud-Maitre Y, Algernon O, Raynaud J. Cytotoxic and antitumoral activity of *Calendula officinalis* extracts. J Pharmazie 1988;43:220-1.
13. Murkute AV, Sahu MS, Mali PY, Rangari VD. Development and Evaluation of Formulations of Microbial Biotransformed Extract of Tobacco Leaves for Hair Growth Potential. J Pharmacognosy Res 2010;2 (5):300-3.
14. Meenavangalapati, Surya Prakash D, Sree Satyanandam. *In-Vitro* Anti-Cancer Studies of Chebulinic acid on colon adenocarcinoma Ht-29 cell lines. Int J Pharm Pharm Sci 2013;5(2):582-3.