

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/350099718>

EFFECT OF NITROGEN AND PANCHAGAVYA ON GROWTH AND YIELD OF BABY CORN (*Zea mays* L.)

Research · January 2020

CITATIONS

0

3 authors, including:



Dr. Shikha Singh

Sam Higginbottom University of Agriculture, Technology and Sciences

21 PUBLICATIONS 1 CITATION

SEE PROFILE

EFFECT OF NITROGEN AND PANCHAGAVYA ON GROWTH AND YIELD OF BABY CORN (*Zea mays* L.)

S.SREETHU* AND SHIKHA SINGH

Department of Agronomy, Naini Agricultural Institute,
SHUATS, Prayagraj, Uttar Pradesh - 211 007
e-mail: itsmesreethu96@gmail.com

KEYWORDS

Baby corn
Nitrogen
Panchagavya

Received on :
20.03.2020

Accepted on :
10.05.2020

*Corresponding
author

ABSTRACT

The study was aimed to analyse the effect of nitrogen and panchagavya on growth and yield of Baby corn (*Zea mays* L.). Among the treatments 60 kg N + 4 sprays of 3% Panchagavya was superior than the rest and recorded the maximum plant height (117.67cm), dry matter accumulation (1466.52 g/m²), leaf area index (8.00), crop growth rate (41.48 g/m²/day), number of cobs per plant (2.14), number of cobs/ha (166296.3), yield with husk (261.39 q/ha), yield without husk (53.71 q/ha) and greenfodder yield (37t/ha) in baby corn. The present investigation revealed that 60 kg N + 4 sprays of 3% Panchagavya would produce maximum growth and yield attributes and can be recommended for baby corn cultivation.

INTRODUCTION

Maize (*Zea mays* L.) is the world's third leading cereal crop following wheat and rice, and has the highest production potential among the cereals. In India, maize is grown in 9.22 M ha area with a production and productivity of 28.72 million tonnes and 3,115 kg/ha respectively contributing 2.53% share over world's production (Directorate of Economics and Statistics 2018, FAO Statistics 2020). Baby corn is a vegetable crop that can potentially improve the economic status of farmers (Das *et al.*, 2008). Baby corn is the young, finger-length de-husked corn young ear of female inflorescence, harvested within 2-3 days of silk emergence but prior to fertilization and is crisp and sweet in taste. (Pandey *et al.*, 2000). It is a profitable crop that allows a diversification of production, aggregation of value, and increased income (Pandey *et al.*, 2002). It is highly remunerative and farmers can get a high return in a short period of 45-60 days.

As baby corn is a high demanding crop which provide more income within a short period, farmers are cultivating those on large scale with huge input of inorganic fertilisers which in turn leads to land degradation. The main idea of this research is to supplement some source of organic formulations in replacement of inorganic fertilizer.

Nitrogen is most deficient primary nutrient in Indian soil and varies from state to state. Nitrogen is considered as one of the most important plant nutrients for growth and development of crop plant. It also plays an important role in synthesis of chlorophyll and amino acids that contribute to the building unit of protein and thus, growth of plants. Nitrogen helps in early establishment of leaf area capable of photosynthesis.

Nitrogen promotes leaf and stem growth rapidly which consequently increase the yield and its quality. (Chouhan *et al.*, 2016). Research substantiates that available nitrogen status increased with increased supply of nitrogen in the form of either fertilizers or organic manures which ultimately increased the productivity of maize. As far as grain yield in cereal is concerned the role of nitrogen is vital. It is therefore, necessary to identify the critical steps associated with nitrogen use efficiency. The nitrogen use efficiency is only about 30-40% in Indian soil.

In addition to nitrogen plants require so many vital elements. Panchagavya is one among the effective methods which can provides macro nutrients, essential micro nutrients, many vitamins, required amino acids, growth promoting substances and beneficial microorganisms for plants well growth. Panchagavya is known to contain bio-fertilizers such as Azospirillum, Azotobactor, Phosphobacteria and Psuedomonas besides Lactobacillus (Yadav and Lourduraj, 2006). Foliar spray of panchagavya provides drought hardiness by acting as a thin oily layer on leaves and stems and thus reduce evaporational loss and also it can play a significant role in providing resistance to pests and diseases, resulting in overall yields (Tharmaraj *et al.*, 2011).

Intensive agricultural practices have resulted in numerous problems like micro nutrient deficiencies, nutrients imbalances, and deterioration of soil health and decline crop yield. No single source of nutrient is capable of supplying plant nutrients in adequate amount and in balanced proportion. Not only this, but also fertilizers are more expensive in developing countries. Therefore, the current trend is to explore the possibility of supplementing chemical fertilizers

with organic and Biofertilizers.(Mishra *et al.*, 2016).Hence the paper deals with the effect of nitrogen and panchagavya on growth and yield of baby corn with the aim to include environmentally safe technology for baby corn production.

MATERIALS AND METHODS

The experiment was conducted during the Zaid season 2019, at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (U.P.) This area is situated on the right side of the Yamuna river by the side of Prayagraj-Rewa road about 12 km from the city. Prayagraj has a subtropical and semi-arid climatic condition, south-eastern part of Uttar Pradesh prevails with both extremes of temperature, *i.e.* winter (3°C) and summer(48°C).The soils were characterized as sandy loam in texture with pH 7.1 analyzed using glass electrode pH meter, electrical conductivity 0.26 dS/m analyzed using conductivity meter (Jackson, 1973), organic carbon 0.36% Walkley and Black method (Nelson and Sommer, 1982), available N 171.48 kg/ha Alkaline potassium permanganate method for nitrogen (Subbiah and Assija, 1956), available P 15.20 kg/ha analyzed by Olsen's method (Nelson and Sommer, 1982), available K 232.50 kg/ha analyzed by Flame photometry (Jackson, 1973).

The experiment was laid out in Randomized Block Design. There were 13 treatments and each replicated thrice. T₁:80 kg N + 60 kg P + 40 kg K (control), T₂:40 kg N + 3 sprays of 3% panchagavya, T₃:40 kg N + 3 sprays of 4% panchagavya, T₄:40 kg N + 4 sprays of 3% panchagavya, T₅:40 kg N + 4 sprays of 4% panchagavya, T₆:60 kg N + 3 sprays of 3% panchagavya, T₇:60kg N + 3 sprays of 4% panchagavya, T₈:60 kg N + 4 sprays of 3% panchagavya, T₉:60 kg N + 4 sprays of 4% panchagavya, T₁₀:80 kg N + 3 sprays of 3% panchagavya, T₁₁:80 kg N + 3 sprays of 4% panchagavya, T₁₂:80 kg N + 4 sprays of 3% panchagavya, T₁₃:80 kg N + 4 sprays of 4% panchagavya. Every treatment contains blanket recommendation of phosphorous and potassium (60 and 40 kg/ha) respectively.

For growth attributes, the height of plant was measured from the base of the plant to the highest point of the uppermost leaf

whose tip is pointing down, Number of green leaves/plant was counted, to calculate leaf area index leaves were collected from the plant and leaf area was measured with leaf area meter was divided by ground area and multiplied with number of leaves, crop growth rate represented as dry weight gained by a unit area of crop in a unit time expressed as g/m²/day.

For yield attributes number of cobs and weight of cobs per plant are counted, the cobs were harvested from net plot area as per treatment and weighed without husk and yield was converted into q/ha and from the net plot after harvesting of cobs left over green plants were cut from ground level which were weighed and mean value was recorded for green fodder estimation was converted to t/ha.

The raw data was subjected to appropriate statistical procedure as suggested by Gomez and Gomez (1984). The data from the experiments were analyzed statistically, wherever treatment differences were found significant, the critical differences were worked out at 5% level of probability (P = 0.05).

RESULTS AND DISCUSSION

Nitrogen and panchagavya levels showed substantial effect on different growth and yield attributes of baby corn. Data regarding growth parameters is presented in Table1. The highest plant height was observed with the application of 60 kg N + 4 spray of 3% Panchagavya (117.67cm) which was significantly superior over control (88.00cm), by 33.7 %. This increment indicates that plants used N during active cell division to form building blocks (protein) for cell elongation. Many research studies have showed that organic formulations release nutrients slowly and may reduce the leaching losses, particularly N (Nevens and Reheul, 2003 and Naveed *et al.*, 2008).The highest number of leaves was observed with 60kg N + 3 sprays of 4% panchagavya (9.53) and there was no significant difference among treatments. Increasing nitrogen level did not significantly affect number of leaves per plant. This was probably attributable to the genetic factor which control number of leaves per plant.This was in accordance with the findings of Eltelib *et al.* (2006). The highest dry matter accumulation was observed in 60 kg N + 4 sprays of 3%

Table 1: Effect of different levels of Nitrogen and Panchagavya on Growth attributes of Babycorn.

| Treatments | Plant height (cm) | Number of leaves (nos.) | Dry matter accumulation (g/m ²) | Leaf Area Index (LAI) | Crop Growth Rate (g/m ² /day) |
|---------------------------------------|-------------------|-------------------------|---|-----------------------|--|
| 80 kg N + 60 kg P + 40 kg K (control) | 88 | 8.36 | 670.3 | 6.59 | 16.29 |
| 40 kg N + 3 sprays of 3% Panchagavya | 89.67 | 8.93 | 640.68 | 6.08 | 17.41 |
| 40 kg N + 3 sprays of 4% Panchagavya | 91.4 | 8.6 | 736.96 | 6.4 | 18.52 |
| 40 kg N + 4 sprays of 3% Panchagavya | 99.87 | 9 | 651.79 | 5.17 | 16.29 |
| 40 kg N + 4 sprays of 4% Panchagavya | 101.6 | 9.13 | 540.69 | 6.48 | 13.33 |
| 60 kg N + 3 sprays of 3% Panchagavya | 103.73 | 8.6 | 592.53 | 4.99 | 15.55 |
| 60 kg N + 3 sprays of 4% Panchagavya | 114.87 | 9.53 | 803.62 | 5.06 | 19.63 |
| 60 kg N + 4 sprays of 3% Panchagavya | 117.67 | 9.27 | 1466.52 | 8 | 41.48 |
| 60 kg N + 4 sprays of 4% Panchagavya | 82.6 | 8.8 | 959.16 | 5.05 | 24.07 |
| 80 kg N + 3 sprays of 3% Panchagavya | 100.47 | 8.36 | 622.16 | 4.74 | 16.29 |
| 80 kg N + 3 sprays of 4% Panchagavya | 92.07 | 9.6 | 888.8 | 5.28 | 23.33 |
| 80 kg N + 4 sprays of 3% Panchagavya | 114.07 | 8.87 | 951.76 | 5.5 | 22.96 |
| 80 kg N + 4 sprays of 4% Panchagavya | 107 | 9.27 | 744.37 | 5.85 | 21.48 |
| SEm(±) | 7.1 | 0.41 | 34.51 | 0.44 | 4.17 |
| CD (P = 0.05) | 20.73 | NS | 100.72 | 1.28 | 12.16 |

Table 2: Effect of different levels of Nitrogen and Panchagavya on Yield Attributes of Baby corn.

| Treatments | No. of cobs Per Plant | No. of cobs Per ha | Yield with husk(q/ha) | Yield without husk(q/ha) | Fodder (t/ha) |
|---------------------------------------|-----------------------|--------------------|-----------------------|--------------------------|---------------|
| 80 kg N + 60 kg P + 40 kg K (control) | 1.01 | 78518.5 | 170.65 | 30.67 | 20.67 |
| 40 kg N + 3 sprays of 3% Panchagavya | 1.24 | 96296.3 | 215.08 | 45.88 | 19 |
| 40 kg N + 3 sprays of 4% Panchagavya | 1.41 | 110000 | 218.9 | 47.28 | 27 |
| 40 kg N + 4 sprays of 3% Panchagavya | 1.4 | 109259.3 | 221.46 | 47.87 | 25.33 |
| 40 kg N + 4 sprays of 4% Panchagavya | 1.4 | 109259.3 | 216 | 46.66 | 21.33 |
| 60 kg N + 3 sprays of 3% Panchagavya | 1.13 | 87777.8 | 191 | 42.23 | 21.67 |
| 60 kg N + 3 sprays of 4% Panchagavya | 1.41 | 110000 | 227.24 | 49.11 | 33 |
| 60 kg N + 4 sprays of 3% Panchagavya | 2.14 | 166296.3 | 261.39 | 53.71 | 37 |
| 60 kg N + 4 sprays of 4% Panchagavya | 0.78 | 60370.4 | 177.62 | 39.58 | 22.67 |
| 80 kg N + 3 sprays of 3% Panchagavya | 1.31 | 101851.9 | 209.91 | 45.4 | 27.33 |
| 80 kg N + 3 sprays of 4% Panchagavya | 1.14 | 88888.9 | 193.08 | 43.12 | 22 |
| 80 kg N + 4 sprays of 3% Panchagavya | 1.72 | 133703.7 | 221.03 | 49.22 | 30.67 |
| 80 kg N + 4 sprays of 4% Panchagavya | 1.61 | 125555.6 | 233.81 | 50.76 | 25.67 |
| SEm(±) | 0.09 | 7008.8 | 9.99 | 2.49 | 1.29 |
| CD (P = 0.05) | 0.26 | 20457.25 | 29.17 | 7.28 | 3.78 |

Panchagavya (1466.52 g/m²) which was significantly superior over rest of the treatments. The probable reason for maximum dry matter might be due to the supply of all micro and macronutrients and growth enzymes present in the panchagavya which favoured rapid cell division and multiplication. . With respect to Panchagavya, ammonia and nitrite oxidizers were found to colonize the leaves and increased the uptake and total N (Papen *et al.*, 2002) and the presence of indole acetic acid (IAA) stimulated the growth of adventitious roots.

The highest leaf area index (LAI) was observed in 60 kg N + 4 sprays of 3% Panchagavya (8.00) which was significantly higher than rest of the treatments. This might be due to the fact that LAI is related to the supply and availability of N to plants which is supplied by the source of foliar applied Panchagavya. Organic manure supply essential nutrient elements to promote vigorous growth and physiological activities in the plant system (Kumar *et al.*, 2018). In the present study CGR was significantly higher for 60 kg N + 4 sprays of 3% Panchagavya. The possible reason might be due to the fact that leaves are the main factors of photosynthesis and dry matter accumulation is expected that the treatment with optimum LAI will have more CGR. Panchagavya increased synthesis of growth promoting substances which is turn helped in increased growth. Similarly findings have been reported by Swaminathan *et al.* (2007) and Choudhary *et al.* (2014)

Yield parameters presented in Table 2 viz. the highest number of cobs/plant (2.14), highest number of cobs/ha (166296.3), maximum yield with husk (261.39 q/ha), maximum yield without husk (53.71 q/ha) and highest green fodder yield (37.00 t/ha) was observed in 60 kg N + 4 sprays of 3% Panchagavya which is significantly superior over rest of the treatments. The higher yield attributes obtained with the effect of hormonal substances present in panchagavya especially cytokinin which plays a role in vegetative plant parts with nutrient partitioning while in reproductive parts, high levels of nutrient mobilization. Foliar application of panchagavya readily supplied nutrients and growth hormones viz. IAA and GA3 present in panchagavya which might have stimulated the production of growth regulators in cell system. Similar results of higher gross and net returns were obtained with the

application of panchagavya by Yadav and Lourduraj (2006) in rice and Somasundaram (2003) in greengram. The significant effect of panchagavya was mainly attributed to its nutrient content, higher biological activity and presence of plant growth promoting substances, which was confirmed by Hazarika *et al.* (2006).

Hence, the present investigation revealed that 60 kg N + 4 sprays of 3% Panchagavya would produce maximum growth and yield attributes and can be recommended for baby corn cultivation.

REFERENCES

- Choudhary, K.M., Patel, M.M., Pagar, R.D. 2014. Effect of foliar application of panchagavya and leaf extracts of endemic plants on groundnut (*Arachis hypogaea* L.). *Agricultural Research Communication Centre*. **37(2)**: 223-226.
- Chouhan, M., Gudadhe, N. N., Kumar, D., Kumawat, A. K. and Kumar, R. 2016. Transplanting dates and Nitrogen Levels Influences on Growth, Yield Attributes, And Yield Of Summer Pearl Millet. *The Bioscan*. **10(3)**: 1295-1298
- Das, S., Ghosh, G., Kaleem, M.D. and Bahadur, V. 2008. Effect of different levels of nitrogen and crop geometry on the growth, yield and quality of baby corn (*Zea mays* L.) CV. 'GOLDEN BABY'. *ISHS Acta Horticulturae 809: International Symposium on the Socio-Economic Impact of Modern Vegetable Production Technology in Tropical Asia*.
- Directorate of Economics and Statistics. 2018. Department of Agriculture and cooperation, Ministry of Agriculture, Government of India.
- Etelib, H.A., Hamad, M.A. and Ali, E.E. 2006. The effect of nitrogen and phosphorus fertilization on growth, yield and quality of forage maize (*Zea mays* L.) *Ind.J. Agron.* **5(3)**: 1812-5379.
- FAO Statistics. 2020. Statistics division, Food and agricultural organisation. "
- Gomez, A. K. and Gomez, A. A. 1984. *Statistical Procedures for Agricultural Research*, 2nd ed. John Wiley and Sons, New York, pp. 180-209.
- Hazarika, U. K., Munda, G. C., Bujorboruah, K. M., Das, A., Patel, D. P., Prasad, K., Rajeshkumar, Danwar, A. S., Tomar, J. M. S., Bordoloi, J., Sharma, M. and Girin, G. 2006. Components of nutrient management. *Nutrient Management Organic Farming*. pp. 15-53.
- Jackson, M. L. 1973. *Soil chemical Analysis* prentice Hall of India.

Pvt. Ltd. New Delhi .p. 498.

Kumar, M.V., Velayutham, A., Kumar, N.S., and Vasanthi, D. 2018. Influence of different organic manures on the growth and yield of Baby corn. *Int. Jo. Adv. Agr. Sc & Tech.* **7(5):** 167-174.

Mishra, D., and Lal, E.P. 2016. Effect of Bio And Chemical Fertilizers on Soil Properties, Yield Attributes and Yield of Baby Corn (*Zea Mays* L.) *The Bioscan.* **11(1):** 411-415.

Naveed, M., Khalid, M., Jones, D. L., Ahmad, R. and Zahir, Z. A. 2008. Relative efficacy of *Pseudomonas* spp., containing ACCdeaminase for improving growth and yield of maize (*Zea mays* L.) in the presence of organic fertilizer. *Pak. J. Bot.* **40(3):** 1243-1251.

Nelson, D. W. and Sommer, L. E. 1982. Total carbon, organic carbon and organic matter in: *Methods of Soil Analysis Part-II.* Page. A.L. (Ed), Am. Soc. Agron. Inc. Soil Sci. Soc. Am. Madism. pp. 539- 577.

Neuens, F. and Reheul, D. 2003. The application of vegetable, fruit and garden waste (VFG) compost in addition to cattle slurry in a silage maize monoculture: nitrogen availability and use. *Europ. J. Agron.* **19:** 189-203.

Pandey, A.K., Prakash, V., Mani, V.P and Singh, R.D. 2000. Effect of rate of nitrogen and time of application on yield and economics of baby corn (*Zea mays* L.). *Indian J. Agronomy.* **45(2):** 338-343.

Pandey, A.K., Mani, V.P., Ved Prakash, Singh, R.D. and Gupta, H.S.

2002. Effect of varieties and plant densities on yield, yield attributes and economics of baby corn (*Zea mays*). *Indian J. Agronomy.* **47(2):** 221-226.

Papen, H., Gebler, A., Zumbusch, E. and Rennenbeg, H. 2002. Chemolitho autotrophic nitrifiers in the phyllosphere of a spruce ecosystem receiving high nitrogen input. *Current Microbiol.* **44:** 56-60.

Somasundaram, E. 2003. Evaluation of organic sources of nutrients and panchagavya spray on the growth and productivity of maize-sunflower-green gram system. Thesis submitted to Tamil Nadu Agricultural University, Coimbatore for award of Ph. D. degree.

Subbiah, B. V. and Assija, G. L. 1956. A rapid procedure for the estimation of available nitrogen in soil. *Current Sci.* **25(8):** 259-260.

Swaminathan C, Swaminathan V, Vijayalakshmi K. 2007. Panchagavya - Boon to organic farming, International Book, Distributing Co., Lucknow.

Tharmaraj, K, Ganesh, P, Kumar. S.R, Kolanjinathan. K. 2011. A Critical Review on Panchagavya– A Boon Plant Growth. *International J. Pharmaceutical & Biological Archives.* **2:**1611-1614.

Yadav, B. K. and Lourduraj, C. A. 2006. Effect of organic manures and panchagavya spray on yield attributes, yield and economics of rice (*Oryza sativa* L.). *Crop Research.* **32(1):** 1-5.



**NATIONAL ENVIRONMENTALISTS
ASSOCIATION**

The National Environmentalists Association is chartered in Ranchi as a nonprofit scientific and educational association of like minded academicians, researchers, scientists from all over the nation for the furtherance and diffusion of knowledge of Life Sciences in general and Environmental Science in particular.

The association not only honours its members but also provides FELLOWSHIP to outstanding contributors to the subject and the society.

Contact :

For Editorial Information

Prof. M. P. Sinha
Vice Chancellor
Sido Kanhu Murmu University
Dumka - 814 110
Jharkhand, INDIA

For information regarding Association :

SECRETARY,
National Environmentalists Association,
D-13, Sai Roofs, 1st Floor,
H. H. Colony,
Ranchi - 834002
Jharkhand, India

E-mails : editor.bioscan@gmail.com
dr.mp.sinha@gmail.com
nat.env.assoc@gmail.com

Cell : 94313-60645; 9572649448

Ph. : 0651-2244071

Website : www.thebioscan.com
: www.neaindia.org

NAAS Rating : 5.26

U.S.A. Office

2827 Videre Dr.,
Wilmington,
DE 19808 We, USA

Type setter Bandana Solutions Facility Management LLP
Published by Aditi Publications, Patliputra, Patna

The Bioscan

An International Quarterly Journal of Life Sciences

ISSN : 0973-7049

Volume 15(2) : 2020

Published as an official organ by

NATIONAL ENVIRONMENTALISTS ASSOCIATION

CONTENTS

Page

A. REVIEW PAPER

1. Seasonal crop yield forecasting-methods, accuracies and limitations: A Review
M. Hanumanthappa., Vignesh., A. T. Krishnamurthy and Rekha Badalingappanavar ————— 149 - 155

B. RESEARCH PAPER

2. Diuron induced phytotoxicity in cotton grown on red and black soils
Md. Mifta Faizullah, T. Ramprakash, T. Anjaiah and M. Madhavi ————— 157 - 160
3. Effect of organic sources of nitrogen fertilization on growth and yield of pearl millet [*Pennisetum glaucum* (L.) R. Br. Emend Stuntz]
Lal Chand Yadav, L. R. Yadav and Saroj Kumari Yadav— 161 - 163
4. Avifaunal diversity of Mallathahalli lake in Bangalore Urban Dt., Karnataka, India
Vidya Padmakumar, C. Silambarasan and Shine P Joseph ————— 165 - 167
5. Evaluation of the performance of french bean (*Phaseolus vulgaris*) varieties under aluminium stress environment
Beauty Borang and Y. K. Sharma ————— 169 - 172
6. Nutrient utilisation by herbaceous flora in cold desert of Himachal Pradesh, India
Poonam and Rajan Bawa ————— 173 - 176
7. IN Vitro evaluation of plant extracts and fungicides on ceratocystis fimbriata (Ellis and Halst.), incitant of pomegranate wilt
Rajat Sharma and Kishore Khosla ————— 177 - 181
8. Zooplankton diversity and physicochemical parameters of Hiran river and a veraval coastal region, Gujarat, India
Pooja M. Kuvadiya, Jatin V. Raval and Agni Vyas ————— 183 - 188
9. Evaluation of bio-efficacy of newer molecules of insecticides against thrips, *Megalurothrips usitatus* in yard long bean, vigna unguiculata sub sp. sesquipedalis
Ramesh. M. Maradi, K. Rajashekharappa and Kiranamaya Pradhan ————— 189 - 192
10. Study on selection parameters and inter-associations of yield, agronomic traits and canopy temperature in relation to drought tolerance in wheat
J. M. Patel., S. K. Patel, C. R. Patel and D. R. Chaudhari ————— 193 - 197
11. Response of different greengram (*Vigna radiata* L. Wilczek) cultivars to varying plant populations
Brijal R. Patel, D. K. Patel, T. V. Reddy, Grishma N. Patel and M. M. Chaudhary ————— 199 - 203
12. Soil nutrient storage under major ecosystems of cold deserts of Himachal Pradesh

The Journal is Currently Abstracted / Indexed in

- Paryavarn Abstract, INDIA
- Indian Science Abstract, INDIA
- Cambridge Science Abstract, U.S.A.

- Zoological Record, U.K.

- Directory of Open Access Journal (DOAJ)

- Chemical Abstract, U. S. A.

- Research BIB

- Indian Science

- Journal Seek

- Scientific Indexing Service (SIS)

Journal is currently rated by

Index Copernicus

Universal Impact Factor

NAAS

DISCLAIMER

The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher/ Association and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher / Association and Editors of the products advertised.

- Poonam and R. Bawa————— 205 - 207
13. Analyzing the impact of organic manures on soil microbial parameters following okra-broccoli cropping pattern
Vishav Vikas and Jag Paul Sharma————— 209- 212
 14. Management of citrus gummosis caused by *Phytophthora* Spp.
H. N. Prajapati and A. H. Barad————— 213- 217
 15. Impact of cropping pattern on biomass productivity and farming community of cold desert agro ecosystem of Himachal Pradesh
Poonam and Rajan Bawa————— 219- 223
 16. Screening of chickpea cultivars against wilt and root rot complex disease of chickpea
Rajkumari Padamini, Kusum Mathur and Amit Trivedi———— 225- 228
 17. Epidemiology and Management of *Colletotrichum gloeosporioides* causing leaf spot of strawberry (*Fragaria X ananassa* Duch.) in Kerala
P. Amrutha and Reshmy Vijayaraghavan————— 229- 235
 18. The first report on the freshwater malacofauna of Newasa tahsil from upper Godavari basin, Maharashtra
G.S.Pande and S.R. Patil————— 237- 242
 19. Effect of nitrogen and panchagavya on growth and yield of Baby corn (*Zea mays* L.)
S .Sreethu and Shikha Singh————— 243- 246
 20. Effect of growth regulators and corm size on growth and corm parameters in gladiolus Cv. malaviya kundan
Saipriya Panigrahi, Anjana Sisodia and Anil K. Singh—— 247- 251
 21. Effect of foliar spray of nutrient and plant growth regulators on growth and yield of green gram (*Vigna radiata* L.)
D.S.V.S .Sruthi., Shikha Singh, Dhananjay Tiwari and P.Kavya————— 253- 256
 22. Multivariate analysis in Blackgram (*Vigna Mungo* (L.) Hepper)
Duddukur Rajasekhar, B. Nandini Priya Yadav, K. Hemalatha ,Gangishetti Ranjithkumar and Gabriyal M. Lal————— 257- 259
 23. Effect of organic manures and natural growth stimulants on the growth and quality characters of Bhendi
M. Ashok Kumar, C. Malarvizhi and K. Dhanasekaran—— 261- 265
 24. Genetic diversity studies in sweet corn (*Zea mays* L. Saccharata.)
A.Thanga Hemavathy————— 267- 269
 25. Assessment of genetic variability parameters in magic population of soybean [*Glycine Max* (L.) Merrill]
Bidush Ranjan Swar, V. Swarna Latha, M. Rajendar Reddy and S. Vanisri————— 271- 274

ZONAL CO-ORDINATORS OF THE ASSOCIATION

- Prof. N. Behera
School of Life Science,
Sambalpur University
- Dr. Nirmal Kumar
ISTAR, Vallabh Vidyanagar,
Anand, Gujarat
- Dr. P. N. Sudha
D. K. M. College for Women,
Vellore
- Prof. S. P. S. Dutta
Dept. of Environmental Science,
Jammu University, Jammu
- Dr. V. Salom Gnana Thanga
Dept. of Env. Scs.,
University of Kerala, Kariavattom
Tiruvananthapuram, Kerala

Publications of the Association

The Bioscan

An International Quarterly Journal of Life Sciences

The Ecoscan

An International Quarterly Journal of
Environmental Sciences

Both the Journals are
online

Both the Journals are
available on Google.com

Websites of the Journals are
www.theecoscan.in
www.thebioscan.in

FEATURES OF ASSOCIATION

- Association is registered under 80G of I.T.
- Prestigious fellowship of the Association (F. N. E. A.) to academicians of the nation.
- Regular annual conference of national and international levels organized by the Association.
- Young Scientist and Senior Scientist award during the conference of the Association.

| THE BIOSCAN : SUBSCRIPTION RATES | | | | |
|---|--------------|----------------|--------------------|--------------------|
| | | India (Rs.) | SAARC Countries | Other Countries |
| Individuals | One Year | 1,000 | 2,000(I:C) | US \$200 |
| | Life Member* | 10,000 | | |
| Institutions | One Year | 3,000 | 6,000(I:C) | US \$400 |
| | Life Member* | 30,000 | | |

*Life Member will receive the journal for 15 years while other benefits will continue whole life

THE BIOSCAN : MEMBERSHIP FORM

Please enter my subscription for the above journal for the year / life member.

Name:

Address:

E-mail:

Payment Rs. : by DD / MD in favour of
National Environmentalists Association payable at Ranchi, No. Dated
..... is enclosed.

NOTE: FOR MEMBERSHIP THE ABOVE INFORMATION CAN BE SENT ON SEPARATE SHEET